

Medication Authorization Form
(Prescription and Non-prescription Medications)



INSTRUCTIONS:

- Section A must be completed by the parent/guardian for **ALL** medication authorization
- Section B must be completed by the physician prescribing the medication
- **Section A and Section B** must be completed for any **long-term medication authorizations** (those lasting longer than 10 days).
- This authorization is valid from the date below through the end of the current school year.
- A separate form must be used for each medication

Section A: To be completed by parent/guardian

Medication authorization for: (First Name) _____ (Last Name) _____
(Child's Name)

Fairfax United Methodist Church Preschool has my permission to administer the following medication:

Medication: _____

Dosage and time to be administered: _____

Special instructions (if any): _____

Child's known allergy/illness: _____

Symptoms: _____

Parent or Guardian's Signature: _____ **Date:** _____

Parent/Guardian First Name: _____ Last Name: _____ Phone: _____

Section B: To be completed by child's physician

The undersigned certifies it is medically necessary for the medication listed below be administered to

(First Name) _____ (Last Name) _____
(Child's Name)

Medication: _____

Dosage and time to be administered: _____

Special instructions (if any): _____

Child's known allergy/illness: _____

Symptoms: _____

Physician's Signature: _____ **Date:** _____

Physician's First Name: _____ Last Name: _____ Phone: _____



Medication Log

Child's Name _____

Medication: _____

| Date/Time (AM/PM) | Administered by (Sign & print) | Dosage | PRN Symptoms exhibited | Side Effects |
|----------------------|-----------------------------------|--------|---------------------------|--------------|
| 1. | | | | |
| 2. | | | | |

Complete section below for Medication Errors
(if this medication was not given as written on the child's written consent form.)

| Date of Error | Details of medication error (include reason error occurred) | Parents notified | Signature (Sign & print) |
|---------------|--|---|--------------------------|
| 1. | | <input type="checkbox"/> Yes <input type="checkbox"/> No | |
| 2. | | <input type="checkbox"/> Yes <input type="checkbox"/> No | |

If an incident was reported to State of Virginia: Incident Number: _____

Date Reported: _____

Name of representative who took report: _____

Phone No. _____

**PLACE
PICTURE
HERE**

Name: _____ D.O.B.: _____

Allergy to: _____

Weight: _____ lbs. Asthma: Yes (higher risk for a severe reaction) No

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____








THEREFORE:

If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for ANY symptoms.

If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:





SEVERE SYMPTOMS

| | | | |
|---|--|---|---|
|  LUNG Shortness of breath, wheezing, repetitive cough |  HEART Pale or bluish skin, faintness, weak pulse, dizziness |  THROAT Tight or hoarse throat, trouble breathing or swallowing |  MOUTH Significant swelling of the tongue or lips |
|  SKIN Many hives over body, widespread redness |  GUT Repetitive vomiting, severe diarrhea |  OTHER Feeling something bad is about to happen, anxiety, confusion | OR A COMBINATION of symptoms from different body areas. |

↓ ↓ ↓

- INJECT EPINEPHRINE IMMEDIATELY.**
- Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS

| | | | |
|--|--|--|--|
|  NOSE Itchy or runny nose, sneezing |  MOUTH Itchy mouth |  SKIN A few hives, mild itch |  GUT Mild nausea or discomfort |
|--|--|--|--|

FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM AREA, FOLLOW THE DIRECTIONS BELOW:

- Antihistamines may be given, if ordered by a healthcare provider.
- Stay with the person; alert emergency contacts.
- Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.15 mg IM 0.3 mg IM


Antihistamine Brand or Generic: _____

Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____


HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case.
2. Pull off red safety guard.
3. Place black end of Auvi-Q against the middle of the outer thigh.
4. Press firmly, and hold in place for 5 seconds.
5. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN® AND EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.




HOW TO USE EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN

1. Remove the epinephrine auto-injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward.
3. With your other hand, remove the blue safety release by pulling straight up.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'.
5. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
6. Remove and massage the injection area for 10 seconds.
7. Call 911 and get emergency medical help right away.



HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALICK®), USP AUTO-INJECTOR, IMPAX LABORATORIES

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip.
3. Grasp the auto-injector in your fist with the red tip pointing downward.
4. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh.
5. Press down hard and hold firmly against the thigh for approximately 10 seconds.
6. Remove and massage the area for 10 seconds.
7. Call 911 and get emergency medical help right away.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____

PHONE: _____

NAME/RELATIONSHIP: _____

PHONE: _____